

1 **ACCEPTED MANUSCRIPT**

2
3 GENETICS IDENTIFICATION OF SEA HOLLY (*Acanthus ilicifolius*) THROUGH DNA
4 BARCODING FROM COASTAL CILACAP, CENTRAL JAVA, INDONESIA

5
6 Harisam T, Haryono FED, Marhaeni B, Amron, Prayogo NA, Sukardi P

7
8 DOI: -

9
10 To appear in : BIOTROPIA Issue

11
12 Received date : 23 July 2018

13 Accepted date : 05 March 2019

14
15 **This manuscript has been accepted for publication in BIOTROPIA journal. It is**
16 **unedited, thus, it will undergo the final copyediting and proofreading process before**
17 **being published in its final form.**

19 **GENETICS IDENTIFICATION OF SEA HOLLY (*Acanthus ilicifolius*) THROUGH**
20 **DNA BARCODING FROM COASTAL CILACAP, CENTRAL JAVA, INDONESIA****

21
22 **Taufan Harisam^{1,*}, Florencius Eko Dwi Haryono¹, Bintang Marhaeni¹, Amron¹,**
23 **Norman Arie Prayogo^{1,2} and Purnama Sukardi¹**

24 ¹Fisheries and Marine Science Faculty, Jenderal Soedirman University, Purwokerto53122,
25 Indonesia

26 ²Centre for Maritime Bioscience Studies, Institute for Research and Community Service,
27 Jenderal Soedirman University, Purwokerto53122, Indonesia

28 *Corresponding author, e-mail: taufan.ltd@gmail.com

29 **This paper was presented at the 2nd Scientific Communication in Fisheries and Marine
30 Sciences (SCiFiMaS 2018), 07-09 May 2018, Purwokerto, Central Java, Indonesia

31
32 Running title: DNA barcoding sea holly in Indonesia

33
34 **ABSTRACT**

35 Information on molecular taxonomy of Sea Holly (*A. ilicifolius*) in coastal Cilacap,
36 Central Java, Indonesia, was very limited. The present study aims to identified Sea Holly (*A.*
37 *ilicifolius*) in coastal Cilacap. This study produced a reference library of molecular for Sea
38 Holly (*A. ilicifolius*). The recently recorded species were utilized for the barcoding
39 investigation. Genetic identification were evaluated by *rbcL* and *matK* gene. The young leaf
40 samples of *A. ilicifolius* was extracted for DNA isolation and amplified with *rbcL* and *matK*
41 primer. The length of *rbcL* gene was 608 bp, and the *matK* gene was 970 bp. The evolutionary
42 history was build using the Neighbor-Joining Method. The barcodes sequences *rbcL* and
43 *matK* analyzed with BLAST and MULTALIN. The sequences were also submitted to NCBI.
44 From this study, we found that genera of *Acanthus* (Acanthaceae) and other genera were
45 clustered in the same clade with high bootstrap value. The result indicated that locus of *rbcL*
46 and *matK* gene cannot be used for species differentiation in *Acanthus*, however, these genes
47 can be used for distinguishing the genus level within Acanthaceae.

48
49 **Keywords:** *A. ilicifolius*, DNA barcode, *matK*, *rbcL*, sea holly

50
51 **INTRODUCTION**

52 Sea holly (*Acanthus ilicifolius*) are important mangrove plant. The plant commonly
53 grows on the wetlands at the river mouth and coastal areas, and it is a true mangrove species
54 (Ragavan *et al.* 2015). It is often distinguished from the related genera by its spiny leaves,
55 spicate terminal inflorescences, two bracteoles and uniform anthers (Duke, 2006). In several
56 countries, this plant functions as an efficacious medicinal plant (Simlai *et al.* 2013; Van *et*
57 *al.* 2008; Kumar *et al.* 2008; Ganesh *et al.* 2008; Paul & Seenivasan 2017).

58 Identification based on characteristics of morphology, such as leaf shape, shape of
59 flowers, branching stems and root shape is extremely prone to error (Sheue *et al.* 2009, Sahu
60 and Kathiresan, 2012), as the morphology of *A. ilicifolius* cant be affected by geography.

Therefore, molecular identification is key in discriminating different species. In this study, we employed DNA barcoding, which uses short fragment of nucleotide sequence for fast, precise species identification (Dong *et al.* 2012; Kress *et al.* 2008; Li *et al.* 2012; Li *et al.* 2015; Vivas *et al.* 2014). DNA barcoding technology is still the ideal method for fast identification due to its convenience and low cost (Xu *et al.* 2017).

Molecular identification using DNA barcoding is often needed to obtain fast, low cost and accurate identification. The DNA barcoding using mitochondrial DNA were employed to identify up to species level. The *rbcL* and *matK* genes from the chloroplast genome were used as the core barcode in the consortium for the Barcode of Life (CBOL Plant Working Group, 2009). The *rbcL* gene is the large subunit of Ribulose-bisphosphate carboxylase gene. The *matK* gene is located with *trnK* gene and encodes the tRNA (Lys) (UUU). Substitution rate of the *matK* gene is highest among the plastid genes (Radulović *et al.* 2010). DNA barcoding using *matK* can discriminate more than 90 % of species in the Orchidaceae but less than 49 % in the nutmeg family (Kress & Erickson 2007; Newmaster *et al.* 2008). In other research of the flora of Canada, genetic identification using *rbcL* and *matK* genes had revealed 93 % success in species identification, this methods can achieve discrimination up to 95 % with the addition of the *trnH-psbA* intergenic spacer (Burgess *et al.* 2011). In this study, we evaluated the efficiency of the *matK* and *rbcL* gene for the identification Sea Holly (*Acanthus ilifolicius*) from the coastal area of Cilacap.

MATERIALS AND METHODS

Collection of the sample

A total of 7 Young leaves individuals samples was collected from the west coast of Cilacap, with geographical latitude of 8°35' S - 8°48' S and longitude of 108°46' E - 109°03' E.

DNA Isolation

Total DNA were extracted from leaf tissue Sea Holly (*A. ilifolicius*) with Cetyltrimethyl ammonium bromide (CTAB) (Sahu *et al.* 2016). The CTAB buffer was prepared from 20 mM EDTA; 1.4 M NaCl; 2 % PVP-30; 1 % β -mercaptoethanol; 10 % SDS and 10 mg/ml proteinase K and mixed with leaf sample. The suspension was incubated at 60 °C for 60 min and after this, suspension was centrifuged at 14,000 rpm for 10 min at room temperature with equal volume of CIAA (24:1). The aqueous phase was transferred to a new tube and precipitated with 0.6 volume of cold isopropanol (-20 °C) and chilled 7.5 M

ammonium acetate followed by storage at -20°C for 1 h. The precipitated DNA was centrifuged at 14,000 rpm for 10 min at 4°C followed by washing with 70 % ethanol. DNA was finally dissolved in TE buffer (10 mM Tris-HCl, 1 mM Na₂EDTA, pH 8.0). The quantity DNA was evaluated using agarose gel electrophoresis and quality of total DNA was evaluated with nanodrop (Thermo Scientific, USA).

PCR and DNA sequencing

The DNA samples were amplified with *rbcL* and *matK* plastid primers *rbcLa*_f: 5'-ATGTCACCACAAACZAGAGACTAAAGC-3', *rbcL*724 _r : 5'-"GTAAAATCAAGTCCACCRCG"-3', *matK*_390-f : 5'- "CGATCTATTCATT CAATATTTC"-3' and *matK*_1326-r: 5'TCTAGCACACGAAAGTCGAAGT-3' (CBOL Plant Working Group, 2009). The total mixture was 50- μl containing 10–20 ng of template DNA, 200 μM of dNTPs, 0.1 μM primers and 1 unit of Taq DNA polymerase (Thermo Scientific, USA). The temperature profile of the PCR cycle for *rbcL* was 94°C for 4 min; 35 cycles of 94°C for 30 s, 55°C for 30 s, 72°C for 1 min; repeated for 35 cycles, and final extension 72°C for 10 minute. For the amplification of the *matK* gene, the temperatures used were 94°C for 1 min; 35 cycles of 94°C for 30 s, 50°C for 40 s, 72°C for 40 s; repeated for 37 cycles, and final extension 72°C for 5 minute. The amplified products were separated by agarose gel (1.2 %) electrophoresis and stained with ethidium bromide (Saddhe *et al.* 2016). PCR was conducted at Research Laboratory Jenderal Soedirman University, Indonesia and sequencing analysis were performed in 1st Base DNA sequencing service, Malaysia.

Phylogenetic analysis

A phylogenetic tree was constructed with <http://www.phylogeny.fr> for identified the genetic similarity and differences. Sample of *rbcL* and *matK* genes was compared to sequences from 8 species, retrieved from NCBI GenBank. Relationship in *rbcL* and *matK* genes was generated with CLUSTAL W, Treeview version 1.5.2. was used to generate scoring method percent and the unrooted tree. The *rbcL* sample sequence compared with *Acanthus spinosus* (MF349678.1), *Acanthus montanus* (L12592.1), *Sclerochiton kirkii* (JX572958.1), *Acanthopsis spatularis* (KF724239.1), *Acanthus ilifolius* (KM255065.1), *Acanthus ilifolius* (KX231351.1), *Acanthus ebracteatus* (KX231352.1), *Acanthus ilifolius* (KP697342.1) (Figure 2). The *matK* sequence compared with *Acanthus ilifolius* (KX231339.1), *Acanthus ilifolius* (KM255080.1), *Acanthus ebracteatus* (KX231340.1), *Acanthus longifolius* (AJ429326.1), *Acanthus spinosus* (MF350143.1), *Acanthus mollis*

(HE967332.1), *Acanthus montanus* (HQ384511.1), *Aphelandra scabra* (JQ586377.1) (Figure 3).

Data analysis

Sequence alignment for rbcL and matK sequences was generated using Multalin V.5.4.1 (<http://multalin.toulouse.inra.fr/multalin/>). All known sea holly sequences were searched using the 'BLASTn' tool against the NCBI database and highest-scoring hit from each query is taken as the mangrove identification (<https://www.ebi.ac.uk/services> and <https://www.ncbi.nlm.nih.gov>). Similarity percentage matrix was calculated based on Clustal 2.1 (www.ebi.ac.uk). The rbcL sample sequence for similarity percentage compare with *Acanthus ilicifolius* (KP697342.1), *Acanthus ilicifolius* (KM255065.1), *Acanthus ilicifolius* (KX231351.1), *Acanthus ebracteatus* (KX231352.1), *Acanthopsis spathularis* (KF724239.1), *Acanthus montanus* (L12592.1), *Sclerochiton kirkii* (JX572958.1), *Acanthus spinosus* (MF349678.1) (Table 1) and the number of different nucleotides compare with *Acanthus ilicifolius* (KP697342.1), *Acanthus ebracteatus* (KX231352.1), *Acanthopsis spathularis* (KF724239.1), *Acanthus montanus* (L12592.1), *Sclerochiton kirkii* (JX572958.1), *Acanthus spinosus* (MF349678.1), *Aphelandra aurantiaca* (MF349506.1), *Aphelandra scabra* (JQ590024.1), *Sclerochiton harveyanus* (JX572957.1), *Crossandra infundibuliformis* (JQ933287.1), *Rhinacanthus nasutus* (KF381120.1), *Aphelandra sinclairiana* (L01884.1), *Avicennia officinalis* (KP697352.1), *Avicennia marina* (KP697350.1), *Ruellia blechum* (GU135168.1), *Stachytarpheta jamaicensis* (JQ618493.1) (Table 2).

The Similarity percentage of matK sample sequence compare with *Acanthus ilicifolius* (KX231339.1), *Acanthus ebracteatus* (KX231340.1), *Acanthus ilicifolius* (KM255080.1), *Acanthus montanus* (HQ384511.1), *Acanthus spinosus* (MF350143.1), *Acanthus longifolius* (AJ429326.1), *Acanthus mollis* (HE967332.1), *Aphelandra scabra* (JQ586377.1) (Table 3) and the number of different nucleotides compare with *Acanthus ilicifolius* (KX231339.1), *Acanthus ebracteatus* (KX231340.1), *Acanthus ilicifolius* (KM255080.1), *Acanthus montanus* (HQ384511.1), *Acanthus spinosus* (MF350143.1), *Acanthus longifolius* (AJ429326.1), *Acanthus mollis* (HE967332.1), *Aphelandra scabra* (JQ586377.1), *Aphelandra aurantiaca* (JQ589891.1), *Aphelandra sinclairiana* (GQ981937.1), *Kudoacanthus albonervosus* (KX526470.1), *Sclerochiton harveyanus*

(JX517343.1), *Sclerochiton kirkii* (JX518192.1), *Proboscidea altheifolia* (MF963699.1) and *Schlegelia parviflora* (AJ429345.1) (Table 4).

RESULTS AND DISCUSSION

Identification of sea holly (*A. ilicifolius*) used *rbcL* and *matK* gene

Currently, morphology-based identification and the declining group of taxonomists result in weak identification of species. This problem can be solved by using of molecular method for mangrove species identification. DNA barcoding is designed to provide accurate, and automated species identifications through the use of molecular species tags based on short, standard gene regions (Fazekas *et al.* 2008; Sadde *et al.* 2016). In this research, we found that the plastid core markers *rbcL* and *matK* cannot be used as DNA barcoding for based assessment of sea holly from coastal Cilacap. We successfully amplified partial *rbcL* and *matK* gene from chloroplast genome of sea holly (Figure 1).




Figure 1. The Amplification of *rbcL* gene (608 bp) and *matK* gene (970 bp) of *Sea Holly* (*A. ilicifolius*) and used 1 kb marker.

Table 1. Similarity percentage of Sea Holly (*A. ilicifolius*) based on *rbcL* gene with *Acanthus ebracteatus*, *Acanthopsis spathularis*, *Acanthus montanus*, *Sclerochiton kirkii* and *Acanthus spinosus* calculated using Clustal 2.1 (www.ebi.ac.uk)

Species	1	2	3	4	5	6	7
KP697342.1_ <i>Acanthus ilicifolius</i>	100	100	100	100	100	100	100
KM255065.1_ <i>Acanthus ilicifolius</i>	100	100	100	100	100	100	100
KX231351.1_ <i>Acanthus ilicifolius</i>	100	100	100	100	100	100	100
KX231352.1_ <i>Acanthus ebracteatus</i>	100	99.29	98.3	98.3	98.3	98.3	98.4
KF724239.1_ <i>Acanthopsis spathularis</i>	98.4	98.1	98.1	98.1	98.1	98.1	98
L12592.1_ <i>Acanthus montanus</i>	98.2	98.2	98.2	98.2	98.2	98.2	98.3

JX572958.1_ <i>Sclerochiton kirkii</i>	98.2	98.2	98.2	98.2	98.2	98.2	98.2
MF349678.1_ <i>Acanthus spinosus</i>	98.0	98.4	98.0	98.0	98.0	98.0	98.0

The DNA sequence of *rbcL* and *matK* gene was perform in BLAST at NCBI to find similar sequence under the same or different genus within family Acanthaceae and calculated using Clustal 2.1 (www.ebi.ac.uk). The most highly similar identity sequences obtained from the GenBank base on *rbcL* gene are *Acanthus ilicifolius* (100%), *Acanthus ebracteatus* (100%), *Acanthopsis spathularis* (98.4%), *Acanthus montanus* (98.2%), *Sclerochiton kirkii* (98.2%), *Acanthus spinosus* (98.0%) (Table 1). The similarity was 98.4% with *Acanthopsis spathularis*, which has 9 nucleotides difference with all *A. ilicifolius*. *Acanthus montanus* has 10 nucleotides difference with all *A. ilicifolius* with 98.02% similarity (Table 2).

Table 2. Number of different nucleotides in *rbcL* gene of Sea Holly (*A. ilicifolius*) and other species within Acanthaceae.

Species	Accession number	Identity (%)	Number of different nucleotides
<i>Acanthus ilicifolius</i>	Sea holly	100	0
<i>Acanthus ebracteatus</i>	KX231352.1	100	0
<i>Acanthopsis spathularis</i>	KF724239.1	98.4	9
<i>Acanthus montanus</i>	L12592.1	98.2	10
<i>Sclerochiton kirkii</i>	JX572958.1	98.2	10
<i>Acanthus spinosus</i>	MF349678.1	98.0	11
<i>Aphelandra aurantiaca</i>	MF349506.1	98.0	11
<i>Aphelandra scabra</i>	JQ590024.1	98.0	11
<i>Sclerochiton harveyanus</i>	JX572957.1	98.0	11
<i>Crossandra infundibuliformis</i>	JQ933287.1	98.0	11
<i>Rhinacanthus nasutus</i>	KF381120.1	97.4	14
<i>Aphelandra sinclairana</i>	L01884.1	97.7	17
<i>Avicennia officinalis</i>	KP697352.1	96.4	20
<i>Avicennia marina</i>	KP697350.1	96.4	20
<i>Ruellia blechum</i>	GU135168.1	95.0	22
<i>Stachytarpheta jamaicensis</i>	JQ618493.1	95.0	22

Table 3. Similarity percentage and of Sea Holly (*A. ilicifolius*) based on *matK* gene with *Acanthus ebracteatus*, *Acanthopsis spathularis*, *Rhinacanthus nasutus* and *Acanthus spinosus*

Species	1	2	3	4	5	6	7
KX231339.1_ <i>Acanthus ilicifolius</i>	99.3	99.3	99.3	99.3	99.3	99.3	99.3
KX231340.1_ <i>Acanthus ebracteatus</i>	99.3	99.3	99.3	99.3	99.3	99.3	99.3
KM255080.1_ <i>Acanthus ilicifolius</i>	97.9	97.9	97.9	97.9	97.9	97.9	97.9
HQ384511.1_ <i>Acanthus montanus</i>	96.5	96.5	96.5	96.5	96.5	96.5	96.5
MF350143.1_ <i>Acanthus spinosus</i>	96.5	96.5	96.5	96.5	96.5	96.5	96.5
AJ429326.1_ <i>Acanthus longifolius</i>	96.4	96.4	96.4	96.4	96.4	96.4	96.4

HE967332.1_ <i>Acanthus mollis</i>	96.3	96.3	96.3	96.3	96.3	96.3	96.3
JQ586377.1_ <i>Aphelandra scabra</i>	94.8	94.8	94.8	94.8	94.8	94.8	94.8

Table 4. Number of different nucleotides in *matK* gene of Sea Holly (*A. ilicifolius*) and other species within Acanthaceae retrieved from GenBank

Species	Accession number	Identity (%)	Number of different nucleotides
<i>Acanthus ilicifolius</i>	KX231339.1	99.3	6
<i>Acanthus ebracteatus</i>	KX231340.1	99.3	6
<i>Acanthus ilicifolius</i>	KM255080.1	98.1	8
<i>Acanthus montanus</i>	HQ384511.1	96.5	19
<i>Acanthus spinosus</i>	MF350143.1	96.5	26
<i>Acanthus longifolius</i>	AJ429326.1	96.4	23
<i>Acanthus mollis</i>	HE967332.1	96.3	27
<i>Aphelandra scabra</i>	JQ586377.1	94.8	40
<i>Aphelandra aurantiaca</i>	JQ589891.1	94.5	41
<i>Aphelandra sinclairiana</i>	GQ981937.1	94.5	42
<i>Kudoacanthus albonervosus</i>	KX526470.1	94.4	47
<i>Sclerochiton harveyanus</i>	JX517343.1	93.8	48
<i>Sclerochiton kirkii</i>	JX518192.1	93.7	49
<i>Proboscidea altheifolia</i>	MF963699.1	91.4	75
<i>Schlegelia parviflora</i>	AJ429345.1	91.3	79

The *MatK* gene showed poor performance at species level identification, Sea Holly has 99.3% similarity with *Acanthus ebracteatus* (Table 3 and Table 4). Species differentiation was relatively weak for *matK* in the research of DNA barcoding of Poaceae (Saadullah *et al.* 2016). Single locus of DNA barcode could not provide an significant level of differentiation. Therefore a combination of 2 loci with 4 available markers is needed to determine the ability to species differentiation (Gonzalez *et al.* 2009). Combination of *rbcL* + *matK* gene for species differentiation in the reseach of tropical tree species in Guiana was lower (<50 %) (Gonzalez *et al.* 2009). Lower discrimination was reported in complex and closest taxa of *Holcoglossum*, *Lysimachia*, *Curcuma* and *Ficus* using *matK* and *rbcL* gene (Xiang *et al.* 2011; Zhang *et al.* 2012; Li *et al.* 2012; Chen *et al.* 2015). Differentiation with 2-loci *rbcL* + *trnH-psbA* and *matK* + ITS had result 100% differences between species (Purushosthman *et al.* 2014). The combination between *rbcL* + *matK* markers results better performance at species and genus level identification (Sadde *et al.* 2016). Another study report only 72 % is species level resolution using combined between *matK* and *rbcL* (Saddhe *et al.* 2017).




Figure 2. Phylogenetic tree of genus *Acanthus*, *Acanthopsis*, and *Sclerochiton* constructed based on likelihood phylogeny of nucleotide sequences of *rbcL* gene.




Figure 3. Phylogenetic tree of genus *Acanthus* and *Aphelandra* constructed based on likelihood phylogeny of nucleotide sequences of *matK* gene.

The results showed that *rbcL* locus could not differentiate Sea Holly from *Acanthus ilicifolius* and *Acanthus ebracteatus*, but can differentiate with *Acanthus montanus*, *Acanthus spinosus* and others genera (*Acanthopsis*, and *Sclerochiton*) (Figure 2). The result from phylogenetic tree also showed that *matK* locus could not differentiate Sea Holly from *Acanthus ebracteatus*, otherwise *matK* locus can separate *Acanthus longifolius*, *Acanthus spinosus*, *Acanthus mollis*, *Acanthus montanus*, and *Aphelandra scabra* (Figure 3). The Phylogenetic tree was built from the highest likelihood in sequence alignment. The number of genetic change built from horizontal dimension of the phylogenetic tree. Value of 0.005 and 0.002 explains the length of the branch representing the number of nucleotide substitution (number of substitution per 100 nucleotide site). There was no indel (insertion and deletion) found in *rbcL* and *matK* gene sequence of Sea Holly (*A. ilicifolius*).




Figure 4. Multi Alignment of *rbcL* gene sequence of *Acanthus ilicifolius*, *Acanthopsis spathularis*, and *Sclerochiton kirkii*




Figure 5. Multi Alignment of *matK* gene sequence of *Acanthus ilicifolius*, *Acanthus montanus* and *Aphelandra scabra*.

The position of different nucleotides in species with the same or a different number of nucleotide difference was not the same in *rbcL* and *matK*. As seen in alignment result amongst *Acanthus ilicifolius*, *Acanthopsis spathularis*, and *Sclerochiton kirkii* (Figure 4) has

11 point of different nucleotide and *Acanthus ilicifolius*, *Acanthus montanus* and *Aphelandra scabra* (Figure 5) has 36 points of different nucleotide. The result of multi alignment of *rbcL* sequences has been little variation compared with multi alignment of *matK* sequences. These results are in line with Saddhe *et al* 2016 and Harisam *et al* 2018, that *matK* gene more sensitive compared with *rbcL* gene.

There are many molecular marker techniques that are used to identify tropical mangrove species. Several studies suggested that one marker alone was not accurate enough for identification (Dong *et al.* 2012). The *rbcL* and *matK* gene can be considered as a barcode for mangrove species. The highest rate of sequencing was observed in *rbcL* (98.7 %), while amplification as well as sequencing rate of *matK*, was 99 %.

DNA barcode was primary essential for identification in mangrove. Genetic diversity assessment is also essential in endemic species, endangered and rarely found. This information was important for mangrove conservation. Species identification in this research was performed using *rbcL* and *matK* sequence. From this results showed that *rbcL* and *matK* sequence genes cannot separate *A. ilicifolius* from *A. ebracteatus*. However another research in terrestrial plants were success identified with *rbcL* and *matK* sequence (Hollingsworth *et al.* 2009; Kress *et al.* 2010; Kuzmina *et al.* 2012; Sadde *et al.* 2016). The species were assigned to their taxa based on two methods, the similarity-based method using BLAST score based single linkage (BLASTClust) and tree-based method (NJ). Sadde *et al.* (2016) report that the combine of *rbcL* and *matK* gene has significant variation and can be used for identification of *Acanthus ilicifolius* and other mangrove species in India. However, considering sea holly has 100% similarity with two other mentioned species, it is suggested that the *rbcL* gene cannot be used to differentiate *Acanthus* species (Table 2). In contrast, Ross and CBOL reported that species identification success rate using *rbcL* seems to be higher, whereas *rbcL* recovery ranged from 90 to 100 % (Ross *et al.* 2008; CBOL Plant Working Group 2009). While 9 nucleotides difference in *rbcL* gene can already place *Acanthopsis spathularis* as different species from *Acanthus ilicifolius*. Ideally, the range of minimum discrimination for discrimination species from 0.0 - 0.27%, or similarity of 99.74 - 100% to place organisms in the same species (Purushothaman *et al.* 2014). Sadde *et al.* (2016) report that *matK* has significant variation and can be used for DNA barcode in *acanthus* family. The present coding regions of *matK* and *rbcL*, which are often recommended in the DNA barcode research only show moderate variability in *Populus* genus and inefficient for use in species differentiation. In this result we can suggest that one marker alone such as *matK* and *rbcL*

gene were not helpful to identification mangroves at the species level Sea Holly (*Acanthus ilicifolius*).

CONCLUSION

Identification of Sea Holly (*Acanthus ilicifolius*) base on *matK* and *rbcL* gene was not successful to differentiate at the species level, only successfully on the genus level. Single locus of DNA barcode could not provide an adequate level of differentiating, therefore the combination of more than two loci is suggested.

ACKNOWLEDGMENTS

This project was funded by BLU (Junior Lecture Research) of Jenderal Soedirman University.

REFERENCES

- Burgess KS, Fazekas AJ, Kesanakurti PR, Graham SW, Husband BC, Newmaster SG, Percy DM, Hajibabaei M, Barrett SCH. 2011. Discriminating plants.
- CBOL Plant Working Group. 2009. A DNA barcode for land plants. PNAS USA 106:12794–12797.
- Chen J, Zhao J, Erickson DL, Xia N, Kress WJ. 2015. Testing DNA barcodes in closely related species of *Curcuma* (Zingiberaceae) from Myanmar and China. Mol Ecol Resour 15:337–348.
- Dereeper A, Guignon V, Blanc G, Audic S, Buffet S, Chevenet F, Dufayard J-F, Guindon S, Lefort V, Lescot M, Claverie JM, Gascuel O. 2008. Phylogeny.fr: robust phylogenetic analysis for the non-specialist. Nucleic Acids Res 36:465-469. <http://dx.doi.org/10.1093/nar/gkn180>
- Dereeper A, Guignon V, Blanc G, Audic S, Buffet S, Chevenet F, Dufayard J-F, Guindon S, Lefort V, Lescot M, Claverie JM, Gascuel O. 2008. Phylogeny.fr: robust phylogenetic analysis for the non-specialist. Nucleic Acids Res 36:465-469. <http://dx.doi.org/10.1093/nar/gkn180>.
- Dong W, Liu J, Yu J, Wang L, Zhou S. 2012. Highly variable chloroplast markers for evaluating plant phylogeny at low taxonomic levels and for DNA barcoding. PLoS ONE 7:e35071. <http://doi:10.1371/journal.pone.0035071>.
- Duke NC. 2006. Australia's mangroves: the Authoritative Guide to Australia's Mangrove Plants. University of Queensland and Norman C. Duke, Brisbane.
- Fazekas AJ, Burgess KS, Kesanakurti PR, Graham SW, Newmaster SG, Husband BC, Percy DM, Hajibabaei M, Barrett SCH. 2008. Multiple multilocus DNA barcodes from the plastid genome discriminate plant species equally well. PLoS ONE 3:e2802.
- Ganesh, S., and J.J. Vennila, 2010. Screening for antimicrobial activity in *Acanthus ilicifolius*. Arch. Appl. Sci. Res., 2: 311-315.

320 Gonzalez MA, Baraloto C, Engel J, Mori SA, Pétronelli P. 2009. Identification of
321 Amazonian Trees with DNA Barcodes. PLoS ONE 4:e7483.

322 Harisam, R.T., Haryono, F.E.D., Marhaeni, B., Prayogo, N.A., Soedibya, P.H.T. 2018.
323 Genetic mutation in mangrove *Acanthus ilifolicius* base on DNA Barcode (rbcL and
324 matK gen) in the different environment change in coastal Cilacap, Central Java,
325 Indonesia. E3S Web of Conferences Vol. 47 no. 05005. <https://doi.org/10.1051/e3sconf/20184705005>.
326

327 Hollingsworth ML, Clark A, Forrest LL. 2009. Selecting barcoding loci for plants: evaluation
328 of seven candidate loci with species level sampling in three divergent groups of land
329 plants. Mol Ecol Resour 9:439–457.

330 Kress WJ, Erickson DL, Swenson NG. 2010. Advances in the use of DNA barcodes to build
331 a community phylogeny for tropical trees in a Puerto Rican forest dynamics plot.
332 PLoS ONE 5:e15409.

333 Kumar KT, Gorain B, Roy DK, Samanta SK, Pal M, Biswas P, Roy, A. Adhikari D.
334 Karmakar D. 2008. Anti-inflammatory activity of *Acanthus ilicifolius*.
335 Ethnopharmacol 120:7-12.

336 Kuzmina ML, Johnson KL, Barron HR, Hebert PDN. 2012. Identification of the vascular
337 plants of Churchill, Manitoba, using a DNA barcode library. BMC Ecol 12:25.

338 Li HQ, Chen JY, Wang S, Xiong SZ. 2012. Evaluation of six candidate DNA barcoding loci
339 in *Ficus* (Moraceae) of China. Mol Ecol Resour 12:783–790.

340 Li X, Yang Y, Henry RJ, Rossetto M, Wang Y, Chen S. 2015. Plant DNA barcoding: from
341 gene to genome. Biol Rev 90:157–166.

342 Newmaster SG, Fazekas AJ, Steeves RAD, Janovec J. 2008. Testing candidate plant barcode
343 regions in the Myristicaceae. Mol Ecol Resour 8:480–490.

344 Paul T, Seenivasan R. 2017. The antioxidant, anticancer and anticoagulant activities of
345 *Acanthus ilicifolius* L. roots and *Lumnitzera racemosa* Willd. leaves, from southeast
346 coast of India. Applied Pharmaceutical Science 7 (03): 081-087.

347 Polidoro BA, Carpenter KE, Collins L, Duke NC, Ellison AM. 2010. The loss of species:
348 Mangrove extinction risk and geographic areas of global concern. PLoS ONE 5:
349 e10095. DOI: 10.1371/journal.pone.0010095.

350 Purushothaman N, Newmaster SG, Ragupathy S, Stalin N, Suresh D, Arunraj DR,
351 Gnanasekaran G, Vassou SL, Narasimhan D, Parani M. 2014. A tiered barcode
352 authentication tool to differentiate medicinal *Cassia* species in India. *Genet Mol Res*
353 13:2959-2968. <http://dx.doi.org/10.4238/2014.April.16.4>.

354 Purushothaman N, Newmaster SG, Ragupathy S, Stalin N, Suresh D, Arunraj DR,
355 Gnanasekaran G, Vassou SL, Narasimhan D, Parani M. 2014. A tiered barcode
356 authentication tool to differentiate medicinal *Cassia* species in India. *Genet Mol Res*
357 13:2959-2968. <http://dx.doi.org/10.4238/2014.April.16.4>.

358 Radulovici AE, Archambault P, Dufresne F. 2010. DNA Barcodes for Marine Biodiversity:
359 Moving Fast Forward?. Diversity 2: 450-472.

360 Ragavan P, Saxena A, Mohan P.M, Jayaraj R.S.C, Ravichandran K. 2015. Taxonomy and
361 distribution of species of the genus *Acanthus* (Acanthaceae) in mangroves of the
362 Andaman and Nicobar Islands, India. J. Biodiversitas. Vol. 16 (2), page 225-237.

363 Ross HA, Murugan S, Li WLS. 2008. Testing the reliability of genetic methods of species
364 identification via simulation. *Syst Biol* 57:216-230.

365 Saadullah, Khan ZUD, Ashfaq M, Zaib UN. 2016. Identification of the grass family
366 (Poaceae) by using the plant DNA barcodes *rbcL* and *matK*. *Biodiversity and*
367 *Environmental Sciences* 8:175-186.

368 Sahu SK, Kathiresan K. 2012. Molecular Markers: an intricate tool for new insights in
369 mangrove genetics. *Int. J. Adv. Biotechnol. Res* 3:847e863.

370 Sahu SK, Reena S, Kandasamy K. 2016. Multi-gene phylogenetic analysis reveals the
371 multiple origin and evolution of mangrove physiological traits through exaptation.
372 *Estuarine, Coastal and Shelf Science* 183:41e51.

373 Sadde AA, Rahul Arvind Jamdade, Kundan Kumar. 2016. Assessment of mangroves from
374 Goa, west coast India using DNA barcode. *SpringerPlus* 5:1554. DOI
375 10.1186/s40064-016-3191-4.

376 Sambrook J, Fritsch EF, Maniatis T. 1989. *Molecular cloning, a laboratory manual*, vol I,
377 Cold Spring Harbor Laboratory Press, New York, pp 5.1–5.17.

378 Sheue C.R, Liu HY, Tsai CC, Rashid SMA, Yong JWH, Yang YP. 2009. On the morphology
379 and molecular basis of segregation of *Ceriops zippeliana* and *C. decandra*
380 (*Rhizophoraceae*) from Asia. *Blumea* 54: 220–227.

381 Simlai, A, A. Roy. 2013. Biological activities and chemical constituents of some mangrove
382 species from Sundarban estuary: An overview. *Pharmacogn Rev* 7: 170-178.

383 Van Kiem P, Quang TH, Huong TT, Nhung le TH, Cuong NX, Van Minh C, Choi EM, Kim
384 YH. 2008. Chemical constituents of *Acanthus ilicifolius* L. and effect on osteoblastic
385 MC3T3E1 cells. *Arch Pharm Res*; 31:823-9.

386 Vivas CV, Ramiris CSM, Anderson AA, Marccus A, Eduardo MN, Cássio VDB, Fernanda
387 AG. 2014. DNA barcoding in Atlantic Forest plants: What is the best marker for
388 Sapotaceae species identification?. *Genetics and Molecular Biology* 37 (4): 662-670.

389 Wang WQ, Wang M. 2007. *The Mangroves of China*. Science Press, Beijing [Chinese].

390 Xiang XG, Hu H, Wang W, Jin XH. 2011. DNA barcoding of the recently evolved genus
391 *Holcoglossum* (Orchidaceae: Aeridinae): a test of DNA barcode candidates. *Mol Ecol*
392 *Resour* 11:1012–1021.

393 Xu, S., He Z., Zhang, Z., Guo, Z., Guo, W., Lyu, H., Li, L., Yang, M., Du, Z., Huang,
394 Y., Zhou, R., Zhong, C., Boufford, D.E., Lerdau, M., Wu, C.i., Duke, N.C., Shi, S.
395 2017. The origin, diversification and adaptation of a major mangrove clade
396 (*Rhizophoreae*) revealed by whole-genome sequencing. *National Science Review*.
397 <https://doi.org/10.1093/nsr/nwx065>.

398 Zhang CY, Wang FY, Yan HF, Hao G, Hu CM, Ge XJ. 2012. Testing DNA barcoding in
399 closely related groups of *Lysimachia* L. (Myrsinaceae). *Mol Ecol Resour* 12:98–108.

400 Widayastuti A, Yani E, Nasution KE, Rochmatino. 2018. Diversity of mangrove vegetation
401 and carbon sink estimation of Segara Anakan Mangrove Forest, Cilacap, Central Java,
402 Indonesia. *Biodiversitas* 19 (1) 246-252.